

Tableau des dérivées

$(x^n)' = n \cdot x^{n-1} \quad \forall x \in \mathbb{R} \quad \forall n \in \mathbb{N} \setminus \{1\}$	$(U^n)' = n \cdot U' \cdot U^{n-1} \quad , \quad U(I) \subseteq \mathbb{R}$
$(\sqrt{x})' = \frac{1}{2\sqrt{x}} \quad \forall x \in]0; +\infty[= \mathbb{R}_+^*$	$(\sqrt{U})' = \frac{U'}{2\sqrt{U}} \quad , \quad U(I) \subseteq]0; +\infty[= \mathbb{R}_+^*$
$\left(\frac{1}{x}\right)' = -\frac{1}{x^2} \quad \forall x \in \mathbb{R}^*$	$\left(\frac{1}{U}\right)' = \frac{-U'}{U^2} \quad , \quad U(I) \subseteq \mathbb{R}^*$
$(e^x)' = e^x \quad \forall x \in \mathbb{R}$	$(e^U)' = U' \cdot e^U \quad , \quad U(I) \subseteq \mathbb{R}$
$(\ln(x))' = \frac{1}{x} \quad \forall x \in]0; +\infty[= \mathbb{R}_+^*$	$(\ln(U))' = \frac{U'}{U} \quad , \quad U(I) \subseteq]0; +\infty[= \mathbb{R}_+^*$
$(\sin(x))' = \cos(x) \quad \forall x \in \mathbb{R}$	$(\sin(U))' = U' \cdot \cos(U) \quad , \quad U(I) \subseteq \mathbb{R}$
$(\cos(x))' = -\sin(x) \quad \forall x \in \mathbb{R}$	$(\cos(U))' = -U' \cdot \sin(U) \quad , \quad U(I) \subseteq \mathbb{R}$
$(\tan(x))' = \frac{1}{\cos^2(x)} = 1 + \tan^2(x)$ $\forall x \in \mathbb{R} - \left\{ \frac{\pi}{2} + k\pi; k \in \mathbb{Z} \right\}$	$(\tan(U))' = \frac{U'}{\cos^2(U)} = \left(1 + \tan^2(U) \right) \cdot U'$ $U(I) \subseteq \mathbb{R} - \left\{ \frac{\pi}{2} + k\pi; k \in \mathbb{Z} \right\}$

Tableau des Primitives

$\int x^{\alpha} .dx = \frac{x^{\alpha+1}}{\alpha+1} + cte ; \alpha \neq -1$	$\int U'.U^{\alpha} dx = \frac{U^{\alpha+1}}{\alpha+1} + cte ; \alpha \neq -1$
$\int \frac{1}{2\sqrt{x}} dx = \sqrt{x} + cte$	$\int \frac{U'}{2\sqrt{U}} dx = \sqrt{U} + cte$
$\int e^x .dx = e^x + cte$	$\int U'.e^U .dx = e^U + cte$
$\int \frac{1}{x} dx = \ln(x) + cte$	$\int \frac{U'}{U} dx = \ln(U) + cte$
$\int \frac{1}{x^2} dx = -\frac{1}{x} + cte$	$\int \frac{U'}{U^2} dx = -\frac{1}{U} + cte$
$\int \cos(x) .dx = \sin(x) + cte$	$\int U'.\cos(U) .dx = \sin(U) + cte$
$\int \sin(x) .dx = -\cos(x) + cte$	$\int U'.\sin(U) .dx = -\cos(U) + cte$
$\int \frac{1}{\cos^2(x)} .dx = \tan(x) + cte$	$\int \frac{U'}{\cos^2(U)} .dx = \tan(U) + cte$
$\int (1 + \tan^2(x)) .dx = \tan(x) + cte$	$\int U'.(1 + \tan^2(U)) .dx = \tan(U) + cte$
$\int \frac{1}{1+x^2} .dx = \arctan(x) + cte$	$\int \frac{U'}{1+U^2} .dx = \arctan(U) + cte$